ureter calculus

ureter calculus refers to the formation of stones within the ureter, the tube that carries urine from the kidneys to the bladder. This condition can lead to significant discomfort, obstructive symptoms, and potential complications if not addressed. In this comprehensive article, we will explore the definition, causes, symptoms, diagnosis, treatment options, and preventive measures related to ureter calculus. By understanding these aspects, individuals can better manage their health and seek appropriate medical advice when necessary.

This article will cover the following topics:

- Understanding Ureter Calculus
- Causes of Ureter Calculus
- Symptoms of Ureter Calculus
- Diagnosis of Ureter Calculus
- Treatment Options for Ureter Calculus
- Prevention of Ureter Calculus

Understanding Ureter Calculus

Ureter calculus, commonly known as ureteral stones, are solid masses that form from crystals in the urine. These stones can vary in size from tiny grains to larger formations that can obstruct the urinary tract. The occurrence of ureteral stones is a common condition, affecting millions of people worldwide. The composition of these stones can differ, with the most prevalent types being calcium oxalate, uric acid, struvite, and cystine stones.

The formation of ureter calculus is often a result of multiple factors, including genetic predisposition, dietary influences, and the body's metabolic processes. When certain substances in the urine become overly concentrated, they can crystallize and form stones. If these stones migrate from the kidneys into the ureters, they can cause significant pain and urinary obstruction.

Causes of Ureter Calculus

The formation of ureter calculus can be attributed to several underlying factors. Understanding these causes is essential for effective prevention and treatment.

Dietary Factors

Certain dietary choices can increase the risk of developing ureteral stones. High intake of oxalate-rich foods, such as spinach, nuts, and chocolate, may contribute to calcium oxalate stone formation. Additionally, excessive consumption of salt can lead to increased calcium in the urine, promoting stone development.

Dehydration

Inadequate fluid intake can result in concentrated urine, which facilitates the crystallization of minerals and the formation of stones. Individuals who do not drink enough water are at a higher risk for developing ureter calculus.

Metabolic Disorders

Metabolic disorders can significantly influence stone formation. Conditions such as hyperparathyroidism and renal tubular acidosis can disrupt the balance of minerals in the body, leading to an increased likelihood of stone formation.

Family History

Genetics play a crucial role in determining an individual's susceptibility to ureter calculus. A family history of kidney stones can suggest an inherited tendency toward stone formation.

Symptoms of Ureter Calculus

Recognizing the symptoms of ureter calculus is vital for timely diagnosis and treatment. Symptoms can vary depending on the size of the stone and its location within the urinary tract.

Severe Pain

One of the hallmark symptoms of ureter calculus is severe pain, often described as a sharp, cramping sensation in the back or side. This pain may radiate to the lower abdomen and groin, commonly referred to as renal colic.

Urinary Symptoms

Individuals may experience changes in their urinary patterns, including increased frequency of urination, urgency to urinate, and difficulty urinating. Blood in the urine, known as hematuria, may also occur.

Nausea and Vomiting

The intense pain associated with ureter calculus can lead to nausea and vomiting. These symptoms often accompany the pain and can exacerbate the individual's discomfort.

Diagnosis of Ureter Calculus

Diagnosing ureter calculus typically involves a combination of medical history, physical examination, and imaging studies.

Medical History and Physical Examination

A healthcare provider will first take a detailed medical history, asking about symptoms, dietary habits, and family history of kidney stones. A physical examination may include palpation of the abdomen and kidneys to assess for tenderness.

Imaging Studies

To confirm the presence of ureter calculus, various imaging techniques may be employed:

- X-rays: Can reveal certain types of stones, particularly calcium-based stones.
- Ultrasound: A non-invasive method that can detect stones in the kidneys

and ureters without radiation.

• CT scans: Highly effective in identifying stones of all types and sizes, providing detailed images of the urinary tract.

Treatment Options for Ureter Calculus

The treatment of ureter calculus depends on the size and type of the stone, as well as the severity of symptoms.

Observation and Conservative Management

For small stones that are not causing significant obstruction or pain, a conservative approach may be recommended. This often includes increased fluid intake and pain management with over-the-counter medications.

Medical Expulsive Therapy

In some cases, medications may be prescribed to help facilitate the passage of stones. Alpha-blockers can relax the muscles in the ureter, making it easier for stones to pass.

Surgical Intervention

Larger stones or those causing severe symptoms may require surgical intervention. Common procedures include:

- **Ureteroscopy:** A minimally invasive procedure where a thin tube is inserted through the urethra and bladder to remove the stone.
- Extracorporeal Shock Wave Lithotripsy (ESWL): A non-invasive procedure that uses sound waves to break stones into smaller fragments that can be passed.
- **Percutaneous Nephrolithotomy:** A surgical procedure to remove larger stones directly from the kidney through a small incision in the back.

Prevention of Ureter Calculus

Preventing the formation of ureter calculus involves lifestyle modifications and dietary changes.

Maintain Adequate Hydration

Drinking plenty of fluids, especially water, can help dilute urine and reduce the concentration of stone-forming substances. It is generally recommended to consume at least 2-3 liters of water daily.

Dietary Adjustments

Making informed dietary choices can significantly impact stone formation. Limiting intake of oxalate-rich foods, reducing salt consumption, and maintaining a balanced diet can help prevent the development of ureter calculus.

Regular Check-ups

Individuals with a history of kidney stones should have regular check-ups with their healthcare provider. Monitoring urinary composition and making necessary adjustments can be crucial in preventing recurrence.

In summary, ureter calculus is a common condition that requires awareness and proactive management. Understanding its causes, symptoms, and treatment options can empower individuals to take charge of their urinary health. Seeking medical advice promptly when experiencing symptoms is essential for effective intervention.

Q: What is ureter calculus?

A: Ureter calculus refers to the formation of stones in the ureter, which can cause obstruction and significant pain. These stones are formed from crystallized substances in the urine.

Q: What are the common types of ureter stones?

A: The most common types of ureter stones include calcium oxalate, uric acid, struvite, and cystine stones, each with varying causes and treatment approaches.

0: How can I tell if I have ureter calculus?

A: Symptoms of ureter calculus may include severe pain in the back or side, changes in urination patterns, nausea, and blood in the urine. If you experience these symptoms, it is essential to seek medical evaluation.

Q: What treatments are available for ureter calculus?

A: Treatment options for ureter calculus range from conservative management and medication for smaller stones to surgical procedures like ureteroscopy and extracorporeal shock wave lithotripsy for larger stones.

Q: Can ureter calculus be prevented?

A: Yes, ureter calculus can often be prevented by maintaining adequate hydration, making dietary adjustments, and having regular health check-ups, especially for those with a history of stones.

0: Is ureter calculus a serious condition?

A: While ureter calculus can be painful and lead to complications such as urinary obstruction or infection, it is typically treatable. Prompt medical attention is important to manage symptoms and prevent complications.

Q: How does diet affect the formation of ureter stones?

A: Certain foods can increase the risk of stone formation. For example, high oxalate foods can contribute to calcium oxalate stones, while excessive salt can elevate calcium levels in urine. A balanced diet is crucial for prevention.

Q: What role does hydration play in preventing ureter calculus?

A: Adequate hydration helps dilute urine, reducing the concentration of minerals that form stones. Drinking plenty of water is one of the simplest and most effective preventive measures against ureter calculus.

Q: Can ureter calculus recur after treatment?

A: Yes, ureter calculus can recur, especially in individuals with a history of stones. Preventive strategies, including dietary changes and hydration, can help reduce the risk of recurrence.

Q: When should I seek medical help for ureter calculus?

A: You should seek medical help if you experience severe pain, difficulty urinating, persistent nausea or vomiting, or visible blood in your urine, as these may indicate complications from ureter calculus.

Ureter Calculus

Find other PDF articles:

 $\underline{http://www.speargroupllc.com/algebra-suggest-006/files?docid=DLH10-5177\&title=intermediate-algebra-pdf-free-download.pdf}$

ureter calculus: The Ureter H. Bergman, 2012-12-06 This volume, focusing on the ureter and the diseases which involve it, is an updated second edition. Many journals and textbooks deal with the physiology, pathology, diagnosis, and therapy of derangements of the urinary tract. In most instances, however, the discussion properly centers on the disease process itself and its primary aspects, with only a tan gential description of effects on the ureter. The editor is therefore correct that the ureter itself should be considered a major organ. Though it has been regarded in the recent past as a simple muscular tube, reacting to stretching or filling by contraction, this simplistic view of ureteral phys iology is changing fast. With expanded knowledge of ureteral physiology, a pharmacology is developing which is becoming useful to the clinician in many ways. One of the most interesting aspects of the ureter is its role in inducing the permanent kidney, the metanephros. Relatively slight displacements in the origin of the ureteral bud result in ectopic ureteral orifices and a wide range of congenital anomalies. An ureteral bud which arises medial to the normal position at the genu of the mesonephric duct results in a lateral, and usually incompetent, ureterovesical junction after the duct is taken up to form a portion ofthe trigone. This appears certainly to be the developmental mechanism which results in primary reflux.

ureter calculus: Studies in the Etiology and Pathology of Pyelitis Ureteritis and Cystitis Cystica Harry Dodge Morse, 1925

ureter calculus: Smith's Textbook of Endourology Arthur D. Smith, 2007 Endourology is a dynamic subspecialty involving closed, controlled manipulation within the genitourinary tract. In the past decade the creative efforts of many urologists, radiologists, and engineers have vastly expanded endoscopic technique, to the great benefit of patients with stones, obstruction, cancer, diverticula, cysts, adrenal disease, varices, and diseases of the bladder. This definitive text addresses every aspect of endourologic procedure including methods of access, operative techniques, complications, and postoperative care. The reader is taken, step-by-step, through cutaneous surgery, ureteroscopy, extracorporeal shock wave lithotripsy, laparoscopy, and lower urinary tract procedures. The principles and function of state-of-the-art endourologic instruments are outlined for each procedure. The authorship reads like a Whoâ \mathfrak{t}^{TM} s Who in endodurology . The breadth and depth of their experience is evident throughout the text.

ureter calculus: The Medical Fortnightly, 1905

ureter calculus: Detailed Diagnoses and Procedures, National Hospital Discharge Survey, 1990 Edmund Graves, 1992

ureter calculus: New York Medical Journal, and Philadelphia Medical Journal, 1922

ureter calculus: Detailed Diagnoses and Surgical Procedures for Patients Discharged from Short-stay Hospitals, United States, 1979 Barbara J. Haupt, 1982

ureter calculus: Kirkes' Handbook of Physiology William Senhouse Kirkes, William Morrant Baker, Vincent Dormer Harris, 1885

ureter calculus: International Record of Medicine and General Practice Clinics Edward Swift Dunster, Frank Pierce Foster, James Bradbridge Hunter, Charles Eucharist de Medicis Sajous, Gregory Stragnell, Henry J. Klaunberg, Félix Martí-Ibáñez, 1908

ureter calculus: National Hospital Discharge Survey, 1988

ureter calculus: CT Urography Stuart G. Silverman, Richard H. Cohan, 2007 Featuring over 500 images, this atlas is the first text on performing and interpreting CT urography. Chapters detail the indications and techniques for CT urography, review the risks of radiation exposure, show how normal urinary tract anatomy and variants appear on CT scans, and demonstrate a wide range of urinary tract abnormalities as they appear on thin-section CT. The final chapter illustrates artifacts and diagnostic pitfalls. Chapters on abnormalities follow a case-based teaching file format. Each case is presented on a two-page spread, with images and succinct discussion of the entity and how CT urography was used to diagnose it.

ureter calculus: Journal of the American Medical Association American Medical Association, 1921 Includes proceedings of the Association, papers read at the annual sessions, and list of current medical literature.

ureter calculus: Index-catalogue of the Library of the Surgeon-General's Office, United States Army Library of the Surgeon-General's Office (U.S.), 1914

ureter calculus: The Journal of the American Medical Association , $1902\,$

ureter calculus: Monthly Cyclopedia and Medical Bulletin , 1900

ureter calculus: The National Home and Hospice Care Survey, ... Summary, 1994

ureter calculus: The Medical Standard, 1909

ureter calculus: Detailed Diagnoses and Procedures, National Hospital Discharge Survey , $1994\,$

ureter calculus: The Urologic and Cutaneous Review, 1920

Related to ureter calculus

Ureter - Wikipedia The ureters are tubes composed of smooth muscle that transport urine from the kidneys to the urinary bladder. In adult humans, the ureters are typically 20–30 centimeters long and 3–4

Ureters: Anatomy, Location, Function & Conditions Ureters are tubes that carry pee from your kidneys to your urinary bladder. Most people have one ureter in each kidney, which connects to the sides of your bladder

Ureteral obstruction - Symptoms and causes - Mayo Clinic Urine leaves the body through another small tube called the urethra. A ureteral obstruction is a blockage in one or both of the tubes (ureters) that carry urine from the kidneys

Ureteral Disorders | Ureters | Ureter Function | MedlinePlus $\,$ Muscles in the ureter walls tighten and relax to force urine down and away from the kidneys. Small amounts of urine flow from the ureters into the bladder about every 10 to 15

Ureter | Definition, Function, & Anatomy | Britannica ureter, duct that transmits urine from the kidney to the bladder. There normally is one ureter for each kidney. Each ureter is a narrow tube that is about 12 inches (30 cm) long.

Ureter: Function and Anatomy - Verywell Health The ureters' role in the process is to carry urine from the kidneys to the bladder. Contractions in the ureter force urine away from the kidneys and into the bladder. The ureters

Anatomy, Abdomen and Pelvis Ureter - StatPearls - NCBI Bookshelf These muscular tubes transport urine from the renal pelvis to the bladder. The ureter's muscular layers are responsible for the peristaltic activity that moves urine from the

Ureters: Anatomy, innervation, blood supply, histology | Kenhub Anatomy and function of the ureters. The ureters are bilateral, muscular, tubular structures, responsible for taking urine from one kidney to the urinary bladder for storage, prior

Ureter Anatomy: Overview, Gross Anatomy, Microscopic Anatomy Each ureter is anatomically and functionally divided into three parts: abdominal, pelvic, and intramural parts **Definition of ureter - NCI Dictionary of Cancer Terms** Anatomy of the male urinary system showing the kidneys, ureters, bladder, and urethra. Urine is made in the renal tubules and collects in the renal pelvis of each kidney. The urine flows from

Ureter - Wikipedia The ureters are tubes composed of smooth muscle that transport urine from the kidneys to the urinary bladder. In adult humans, the ureters are typically 20–30 centimeters long and 3–4

Ureters: Anatomy, Location, Function & Conditions Ureters are tubes that carry pee from your kidneys to your urinary bladder. Most people have one ureter in each kidney, which connects to the sides of your bladder

Ureteral obstruction - Symptoms and causes - Mayo Clinic Urine leaves the body through another small tube called the urethra. A ureteral obstruction is a blockage in one or both of the tubes (ureters) that carry urine from the kidneys

Ureteral Disorders | Ureters | Ureter Function | MedlinePlus Muscles in the ureter walls tighten and relax to force urine down and away from the kidneys. Small amounts of urine flow from the ureters into the bladder about every 10 to 15

Ureter | Definition, Function, & Anatomy | Britannica ureter, duct that transmits urine from the kidney to the bladder. There normally is one ureter for each kidney. Each ureter is a narrow tube that is about 12 inches (30 cm) long. A

Ureter: Function and Anatomy - Verywell Health The ureters' role in the process is to carry urine from the kidneys to the bladder. Contractions in the ureter force urine away from the kidneys and into the bladder. The ureters

Anatomy, Abdomen and Pelvis Ureter - StatPearls - NCBI Bookshelf These muscular tubes transport urine from the renal pelvis to the bladder. The ureter's muscular layers are responsible for the peristaltic activity that moves urine from the

Ureters: Anatomy, innervation, blood supply, histology | Kenhub Anatomy and function of the ureters. The ureters are bilateral, muscular, tubular structures, responsible for taking urine from one kidney to the urinary bladder for storage, prior

Ureter Anatomy: Overview, Gross Anatomy, Microscopic Anatomy Each ureter is anatomically and functionally divided into three parts: abdominal, pelvic, and intramural parts **Definition of ureter - NCI Dictionary of Cancer Terms** Anatomy of the male urinary system showing the kidneys, ureters, bladder, and urethra. Urine is made in the renal tubules and collects in the renal pelvis of each kidney. The urine flows from

Related to ureter calculus

Ureteric Calculi Research (News Medical2y) Ureteric calculi are stones in the ureters, which are the narrow muscular tubes that are responsible for the drainage of urine from the kidneys to the bladder. These stones are mostly made up of

Ureteric Calculi Research (News Medical2y) Ureteric calculi are stones in the ureters, which are the narrow muscular tubes that are responsible for the drainage of urine from the kidneys to the bladder. These stones are mostly made up of

Ureteric Calculus (Medindia12y) About - Stone in the ureter(tube connecting Kidney with bladder). Stones are produced in the kidney and one or many stones slip into the ureter. Diagnosis - Usually **Ureteric Calculus** (Medindia12y) About - Stone in the ureter(tube connecting Kidney with bladder).

Stones are produced in the kidney and one or many stones slip into the ureter. Diagnosis - Usually Alfuzosin Stone Expulsion Therapy for Distal Ureteral Calculi: A Double-Blind, Placebo Controlled Study: From F1000 (Medscape17y) Changes Clinical Practice: All patients with distal stones treated medically should be treated with alpha-blockers in order to benefit from this enhancement of quality of life. These findings will

Alfuzosin Stone Expulsion Therapy for Distal Ureteral Calculi: A Double-Blind, Placebo Controlled Study: From F1000 (Medscape17y) Changes Clinical Practice: All patients with distal stones treated medically should be treated with alpha-blockers in order to benefit from this enhancement of quality of life. These findings will

MET for distal ureteral calculi: not all stones are created equal (Nature7y) Results of two large randomized controlled trials have challenged the use of medical expulsive therapy for the management of ureteral calculi. A new trial now shows that tamsulosin is effective in

MET for distal ureteral calculi: not all stones are created equal (Nature7y) Results of two large randomized controlled trials have challenged the use of medical expulsive therapy for the management of ureteral calculi. A new trial now shows that tamsulosin is effective in

Urolithiasis: The Influence of Stone Size on Management (Medscape3y) Background A 37-year-old female presented with pain in the right lower lateral abdomen. She had experienced a ureteral stone in the past. Investigations Serum electrolyte levels, blood urea nitrogen,

Urolithiasis: The Influence of Stone Size on Management (Medscape3y) Background A 37-year-old female presented with pain in the right lower lateral abdomen. She had experienced a ureteral stone in the past. Investigations Serum electrolyte levels, blood urea nitrogen,

New Guidelines On Ureteral Stone Management Released (Science Daily17y) A new Guideline for the Management of Ureteral Calculi has been released. One of the biggest changes in this document is the shift in indications for ureteroscopy. The American Urological

New Guidelines On Ureteral Stone Management Released (Science Daily17y) A new Guideline for the Management of Ureteral Calculi has been released. One of the biggest changes in this document is the shift in indications for ureteroscopy. The American Urological

A case of staghorn stones in a kidney with an ileal ureter treated by percutaneous **nephrolithotomy** (Nature17y) Background A 59-year-old man was admitted to hospital for investigation of a 1-year history of intermittent hematuria. He had undergone ileal ureteral replacement for left renal stones 36 years

A case of staghorn stones in a kidney with an ileal ureter treated by percutaneous nephrolithotomy (Nature17y) Background A 59-year-old man was admitted to hospital for investigation of a 1-year history of intermittent hematuria. He had undergone ileal ureteral replacement for left renal stones 36 years

Spontaneous Extravasation from the Ureter Secondary to Aneurysm of the Abdominal Aorta (The New England Journal of Medicine10mon) RECENTLY the 27th case of spontaneous or nontraumatic extravasation from the ureter was reported. The purpose of the present communication is to present a case secondary to an arteriosclerotic

Spontaneous Extravasation from the Ureter Secondary to Aneurysm of the Abdominal Aorta (The New England Journal of Medicine10mon) RECENTLY the 27th case of spontaneous or nontraumatic extravasation from the ureter was reported. The purpose of the present communication is to present a case secondary to an arteriosclerotic

Ureteric Calculi (News Medical6y) Medically known as ureterolithiasis, ureteral stone disease is characterized by the formation of calculi (i.e. stones) in the kidneys and obstructing the ureters. The ureters are the slender muscular

Ureteric Calculi (News Medical6y) Medically known as ureterolithiasis, ureteral stone disease is characterized by the formation of calculi (i.e. stones) in the kidneys and obstructing the ureters. The ureters are the slender muscular

Back to Home: http://www.speargroupllc.com